

From: Horevitz, E. & Manoleas, P. (in press). Training needs of professional social workers in integrated behavioral health care settings. *Social Work in Health Care*.

	Heard of?		Learned At-School		Learned On-Job		Use on Job?		Important?	
	n	%	n	%	n	%	n	%	n	%
Functional Assessment	65	77	11	13	27	32	39	46	35	42
Warm Handoff	66	79	7	8	50	60	60	71	57	68
Behavioral Activation	40	48	8	10	14	17	33	40	31	37
Motivational Interviewing*	76	91	22	26	13	17	69	82	63	76
Problem Solving Treatment	50	60	8	10	14	17	40	48	36	43
CBT-PC	71	85	18	21	24	29	56	67	50	60
Relaxation training**	75	89	15	18	17	20	69	82	52	74
Team-based care	71	84	14	17	54	64	68	81	68	81
Chronic Illness (knowledge of)	72	86	9	10	54	64	64	76	70	83
Psychotropic Medication	77	92	14	17	45	54	76	91	72	86
AOD	57	68	9	11	25	30	33	39	29	35
Psychoeducation	72	86	25	30	40	48	70	83	68	81
Curbside Consultation	61	72	2	2	54	64	57	68	57	68
Stepped Care	37	44	3	4	16	19	16	50	16	19
Family Systems	77	92	65	77	5	6	73	87	64	76
Case Management	77	92	31	37	42	50	64	76	59	70
Cultural Competence	76	91	46	56	14	17	74	88	73	87
Standardized Outcome Measures	65	77	8	10	44	52	53	63	48	57
PCMH	57	68	2	2	45	54	50	60	52	44

(Important = very important + important).

*The majority (33% n = 28) learned MI at a training;

**the majority (25% n=21) learned relaxation training via self-teaching.